

Diagnosis and management of autism in adults

Valkanova V, Rhodes F, Allan CL. Diagnosis and management of autism in adults.
Practitioner 2013; 257 (1761):13-16

Dr Vyara Valkanova
BSc (Hons) MD
Associate Clinical Fellow, Department of Psychiatry

Ms Fiona Rhodes
BA (Hons)
Fifth Year Medical Student, Jesus College

Dr Charlotte L Allan
MBChB (Hons) MRCPsych
Clinical Research Fellow, Department of Psychiatry
University of Oxford, Oxford, UK



Practitioner
Medical Publishing Ltd

Diagnosis and management of autism in adults

AUTHORS

Dr Vyara Valkanova

BSc (Hons) MD
Associate Clinical Fellow,
Department of Psychiatry

Ms Fiona Rhodes


BA (Hons)
Fifth Year Medical
Student, Jesus College

Dr Charlotte L Allan

MBChB (Hons)
MRCPsych
Clinical Research Fellow,
Department of Psychiatry

University of Oxford,
Oxford, UK

FIGURE 1
Psychosocial
interventions for
autism in adults



What are the symptoms and signs?



ADULTS WITH AUTISM CAN BE SIGNIFICANTLY DISADVANTAGED IN MANY ASPECTS OF LIFE

including employment, social relationships, physical and mental health.¹ Recognition of the importance of identifying, assessing and managing adults with autism is increasing, particularly with the publication of recent NICE guidelines.² It is vital for adults of all ages, including older people,³ to be accurately diagnosed in order to receive appropriate management.

Autism affects 1.1% of the adult population.⁴ The spectrum of symptoms is wide. Some individuals have above average intelligence and are fully independent, while others have limited independence because of a learning disability.

Developmental delay is a core feature, and autism is usually diagnosed in childhood.⁵ However, the diverse

How should diagnosis be confirmed?

clinical spectrum makes early diagnosis difficult, especially if good language and cognitive skills help the child to develop coping strategies. Hence high-functioning individuals with autism, Asperger's syndrome, may remain undiagnosed until adulthood.⁶

SYMPTOMS AND SIGNS

GPs should suspect autism if patients present with the signs and symptoms listed in table 1, p14.

Diagnostic complexity is significantly increased when autism coexists with other mental disorders. Learning disability is the most common, and is present in up to 60% of people with autism,⁷ however, depression, social anxiety, obsessive compulsive disorder and attention deficit hyperactivity disorder (ADHD) may also be found.

Depression and anxiety may occur due to the psychological and social

What are the recommended interventions?

consequences of autism, while common aetiological factors relating to neurodevelopmental disorders, may predispose to ADHD.

Autism is a lifelong condition characterised by difficulties in two core dimensions:

- Problems in social communication
- Strongly repetitive behaviour, resistance to change or restricted interests

People with autism show symptoms from both core dimensions and the difficulties in these two domains can be manifest in multiple ways, see table 2, p14.

ASSESSMENT

Careful consideration of core signs and symptoms is the first step in assessment see table 1, p14. It may be necessary to make adjustments to the physical environment, and to the duration and pacing of the assessment.²

A collateral history should identify »

early developmental and behavioural problems in different settings i.e. at home, in education or employment.² Sensory and gastrointestinal problems are very common, and should be asked about. Up to 90% of adults with autism without a learning disability are hyper- or hyposensitive to sound, light, colour, smell or taste.⁸ Almost half of adults with autism and a learning disability suffer from gastrointestinal disorders.⁹

Further investigations help to determine which patients should

be referred to specialist services for a comprehensive assessment. The Autism-Spectrum Quotient (AQ-10) is a 10-item questionnaire for people with suspected autism.¹⁰ The advantage of using this in primary care is that it provides a time-efficient, structured way of ascertaining key symptoms and clearly signals those who should be referred for further assessment. It is freely available online (<http://guidance.nice.org.uk/CG142/AQ10Test/pdf/English>).

Biological tests e.g. hearing, sight

and electroencephalography, genetic tests and neuroimaging should not be used during the initial assessment, and are rarely needed as part of a comprehensive assessment.²

CONFIRMING DIAGNOSIS

Patients should be referred if autism is suspected clinically. The AQ-10 provides useful additional information to support referral; those who score above six should be referred.¹⁰ The AQ-10 requires a patient to be able to read and write, and therefore is not appropriate for those with a moderate or severe learning disability. Where this is the case, referral should be based on a brief assessment only: if a patient has two or more behaviours suggestive of autism, see table 3, below, then referral is appropriate.²

‘Patients should be referred if autism is suspected clinically’

A diagnosis of autism should be confirmed by a specialist multidisciplinary team.² The assessment should include a detailed developmental history and information about current and past behaviour, see table 4, opposite. This information will usually require a collateral history from someone who knows the patient well. Careful observation is needed to assess the core signs of autism, particularly patterns of social interaction.

The use of formal assessment tools which cover multiple domains including social interaction, communication, and imaginative use of materials, may aid the diagnosis, particularly in complex cases, see table 5, opposite.

However, all assessment tools have major limitations and should only be used in conjunction with a detailed clinical history.¹¹

The results of the assessment should be communicated clearly to the patient, ideally with the support of a friend, family member or carer. If a diagnosis of autism is made, clinicians should do a risk assessment and formulate risk and crisis management plans. These should include details of the roles and responsibilities of both the specialist team and primary care team in managing crisis situations.

Table 1

Signs and symptoms that prompt assessment for autism in adults²

One or more of:	AND	One or more of:
● Persistent difficulties in social interaction ● Persistent difficulties in social communication ● Stereotypic behaviours, resistance to change or restricted interests		● Problems obtaining or sustaining employment or education ● Difficulties initiating or sustaining social relationships ● Contact with mental health or learning disability services ● Neurodevelopmental condition

Table 2

Core symptoms of autism

Social communication difficulties	Strongly repetitive behaviour/resistance to change or restricted interests
● Atypical eye contact ● Intrusion into others' personal space ● Difficulties with two-way interaction and interpreting non-verbal communication (including reduced empathy) ● Tendency to take language literally ● Social naivety and vulnerability to exploitation ● Reduced interest in socialising and difficulties interacting in social groups ● Loneliness and social anxiety (and risk of depression) ● Social anxiety	● Preference for repetition and routine ● Need for sameness (eating the same foods, wearing the same clothes) and anxiety at any change ● Preference for predictability and predictable events (watching washing machines spinning) ● Strict order and precision ● Need for clarity, strict order and precision ● Attention to small 'irrelevant' details ● Narrow deep interests, 'fixated' interests

Table 3

Behaviours indicating autism in adults with a moderate or severe learning disability

- Problems with social interaction including: being aloof, indifferent or unusual, interacting with others only when necessary, difficulty with two-way interaction
- Lack of responsiveness to other people
- Difficulty adapting behaviour to different social situations
- Limited expression of empathy
- Repetitive activities (spinning, hand flapping, rocking) which become more prominent at times of stress or are used as a way of expressing emotion

Adapted from NICE CG142. Autism in adults²

Table 4**Specialist assessment of autism**

- Developmental history
- Social interactions, stereotypic behaviour, sensitivity to sensory stimulus, resistance to change, restricted interests
- Current and past behaviour problems
- Functional capability at home and work
- Past psychiatric history
- Past medical history (to include enquiry about neurodevelopmental conditions and epilepsy)

Adapted from NICE CG142. Autism in adults ²

Table 5**Specialist assessment tools for diagnosing autism**

- Adult Asperger Assessment (AAA)¹⁵
- Autism Diagnostic Interview* – Revised (ADI-R)¹⁶
- Autism Diagnostic Observation Schedule* – Generic (ADOS-G)¹⁷
- Asperger's syndrome (and high-functioning autism) Diagnostic Interview (ASDI)¹⁸

* suitable for people with a learning disability

The role of the GP in a crisis situation will vary depending on the needs of the individual, and may include, acting as the first point of contact (particularly out of hours), assisting the patient in contacting the specialist team, and, if appropriate, ensuring prompt prescription of medications.

If there is disagreement about the diagnosis, a GP could consider obtaining a second opinion. This may be useful, if a patient has complex physical or psychiatric comorbidities, or if local teams lack sufficient expertise in adults with autism.

MANAGEMENT**Psychosocial interventions**

For adults with autism a group-based or individual social learning programme to improve social interaction is recommended.² This should include modelling, decision making, strategies for dealing with socially difficult situations and peer or individual feedback. Training in life skills is recommended for adults with all ranges of intellectual ability, who need help with activities of daily living. This consists of structured training programmes based on behavioural principles. For higher functioning individuals structured leisure activity programmes, and anger management may be beneficial, see figure 1, p13.

Adults with autism have high levels of unemployment,¹² and employment programmes have been successfully

Table 6**Biomedical interventions that are not recommended for the core symptoms of autism**

- Exclusion diets (gluten or caesin-free and ketogenic diets)
- Vitamins, minerals and dietary supplements
- Cholinesterase inhibitors
- Anticonvulsants
- Antipsychotics
- Antidepressants
- Hormonal therapies
- Hyperbaric oxygen therapy

used to support people in obtaining or maintaining employment. Programmes typically include assistance with written applications, interviews, identification of appropriate roles and employer education and support. Prospects, run by the National Autistic Society, placed 68% of clients in work, the majority on permanent contracts, with high client and employer satisfaction and significant job retention at eight year follow-up.¹³

Biological treatments

Further research into the use of antidepressants in managing the core symptoms of autism is needed, and their use is not currently recommended by NICE.²

There is promising trial evidence that fluoxetine significantly improves repetitive behaviours compared with placebo, and is well tolerated.¹⁴ However, interpretation of this study and trials of other SSRIs are limited by small sample sizes and use of scales not validated in autism.

Biomedical interventions that are not recommended for the core symptoms of autism are listed in table 6, above.

CHALLENGING BEHAVIOUR

Before considering any psychosocial or pharmacological intervention for challenging behaviour, an assessment should be made of factors that may be triggering or maintaining the behaviour, including any physical symptoms, coexisting mental disorders and the physical or social environment.²

First-line treatment should focus on psychosocial interventions relevant to unmet needs, triggers and reinforcements of the challenging behaviour, and a complete evaluation of the behaviour, the individual and their environment.²

Intervention strategies should clearly identify target behaviour, desired outcomes, and a timescale for assessment of efficacy and modification. Any intervention should be focused on overall quality of the patient's life.

Second-line treatments may include medication such as antipsychotic drugs. These should be used in conjunction with a psychosocial intervention.² Antipsychotics should not be used first line, unless the patient's behaviour is so dangerous that it precludes the use of psychosocial or other interventions. Antipsychotics should be prescribed by a specialist and should be discontinued if there is no clinically important response by six weeks. Anticonvulsants should not be used routinely for challenging behaviour.

COEXISTING DISORDERS

NICE recommendations are available for coexisting mental disorders,² however, it may be helpful to seek advice from a specialist autism team to discuss how to adapt management.

When considering any pharmacological treatment, low doses should be used and attention should be given to the potential for greater sensitivity to any side effects and idiosyncratic responses.

'Families of adults with autism should be offered an assessment of their own needs'

PRIMARY CARE TEAM SUPPORT

An understanding of autism and the difficulties it poses is central to offering appropriate support to patients and their families.

A significant proportion of adults with autism experience social and economic exclusion, and are unable to access the support they need to live independently.² They are also at increased risk of other mental and physical disorders, and are less likely to report these.

People with autism, and if appropriate their carers, should have the opportunity to be involved in making informed decisions about their care, in partnership with healthcare professionals.² It is helpful to consider the impact of the »

key points

SELECTED BY

Dr Peter Saul

GP, Wrexham and Associate GP Dean for North Wales

Autism affects 1.1% of the adult population. The spectrum of symptoms is wide; some individuals have above average intelligence and are fully independent, while others have limited independence because of a learning disability. Developmental delay is a core feature, and autism is usually diagnosed in childhood. High-functioning individuals with autism, Asperger's syndrome, may remain undiagnosed until adulthood.

Autism is a life-long condition characterised by problems in two core dimensions: difficulties with social communication and strongly repetitive behaviour, resistance to change or restricted interests. The history should identify early developmental and behavioural problems in different settings e.g. at home, in education or employment. Sensory and GI problems are very common, and should be asked about. Up to 90% of adults with autism without a learning disability are hyper- or hyposensitive to sound, light, colour, smell or taste. Almost half of adults with autism and a learning disability suffer from GI disorders.

The Autism-Spectrum Quotient (AQ-10) is a 10-item questionnaire for people with suspected autism. The advantage of using this in primary care is that it provides a time-efficient, structured way of ascertaining key symptoms and clearly signals those who should be referred for further assessment. Biological tests should not be used during the initial assessment. Patients should be referred if autism is suspected clinically and a diagnosis of autism should be confirmed by a specialist multidisciplinary team. If a diagnosis of autism is made, clinicians should do a risk assessment and formulate risk and crisis management plans. These should include details of the roles and responsibilities of both the specialist team and primary care team in managing crisis situations.

For adults with autism a group-based or an individual learning programme to improve social interaction is recommended. Adults with autism have high rates of unemployment, and employment programmes have been successfully used to support people. There is promising trial evidence that fluoxetine significantly improves repetitive behaviours, and is well tolerated. Before considering any psychosocial or pharmacological intervention for challenging behaviour, an assessment should be made of factors that may be triggering or maintaining the behaviour, including any physical symptoms, coexisting mental disorders and the physical or social environment. First-line treatment should focus on psychosocial interventions relevant to unmet needs, triggers and reinforcements of the challenging behaviour, and a complete evaluation of the behaviour, the individual and their environment.

People with autism, and if appropriate, their carers, should have the opportunity to be involved in making informed decisions about their care, in partnership with healthcare professionals. It is helpful to consider the impact of the practice environment, including noise, proximity of others and time pressure.

practice environment, including noise, proximity of others and time pressure. It is likely to take longer to establish a rapport; so it may help to use factual language, visual aids, written information and to avoid abstract terms.

Families of adults with autism provide much of the care and support, and are often crucial in the success of interventions. They should be offered an assessment of their own needs, including psychological, social and emotional support, respite care, emergency plans and planning of future care,² and signposting to relevant organisations.

REFERENCES

- 1 Howlin P, Moss P. Adults with autism spectrum disorders. *Can J Psychiatry* 2012;57(5):275-83
- 2 National Institute for Health and Clinical Excellence. CG142. Autism: recognition, referral, diagnosis and management of adults on the autism spectrum. NICE. London. 2012
- 3 Mukaetova-Ladinska EB, Perry E, Baron M, Povey C. Ageing in people with autistic spectrum disorder. *Int J Geriatr Psychiatry* 2012;27(2):109-18. doi: 10.1002/gps.2711. Epub 2011/05/04
- 4 Brugha T, Cooper SA, McManus S. Estimating the prevalence of autism spectrum conditions in adults: Extending the 2007 Adult Psychiatric Morbidity Survey. The Health and Social Care Information Centre. London. 2012
- 5 Frith U. Autism: Explaining the Enigma. Blackwell. Oxford. 1989
- 6 Attwood T. Asperger's Syndrome: A Guide for Parents and Professionals. Jessica Kingsley. London. 1997
- 7 Baird G, Simonoff E, Pickles A et al. Prevalence of disorders of the autism spectrum in a population cohort of children in South Thames: the Special Needs and Autism Project (SNAP). *Lancet* 2006;368(9531):210-5. Epub 2006/07/18
- 8 Crane L, Goddard L, Pring L. Sensory processing in adults with autism spectrum disorders. *Autism* 2009;13(3):215-28
- 9 Galli-Carminati G, Chauvet I, Deriaz N. Prevalence of gastrointestinal disorders in adult clients with pervasive developmental disorders. *J Intellect Disabil Res* 2006;50(Pt 10):711-8
- 10 Allison C, Auyeung B, Baron-Cohen S. Toward brief "Red Flags" for autism screening: The Short Autism Spectrum Quotient and the Short Quantitative Checklist for Autism in toddlers in 1,000 cases and 3,000 controls [corrected]. *J Am Acad Child Adolesc Psychiatry* 2012;51(2):202-12 e7. Epub 2012/01/24
- 11 Stoesz BM, Montgomery JM, Smart SL, Hellsten LA. Review of five instruments for the assessment of Asperger's disorder in adults. *Clin Neuropsychol* 2011;25(3):376-401. Epub 2011/03/11
- 12 Barnard J, Harvey V, Potter D, Prior A. Ignored or Ineligible? The Reality for Adults with Autism Spectrum Disorders. National Autistic Society. London. 2001
- 13 Howlin P, Alcock J, Burkin C. An 8 year follow-up of a specialist supported employment service for high-ability adults with autism or Asperger syndrome. *Autism* 2005;9(5):533-49. Epub 2005/11/17
- 14 Hollander E, Soorya L, Chaplin W et al. A double-blind placebo-controlled trial of fluoxetine for repetitive behaviors and global severity in adult autism spectrum disorders. *Am J Psychiatry* 2012;169(3):292-9. Epub 2011/12/24
- 15 Baron-Cohen S, Wheelwright S, Robinson J, Woodbury-Smith M. The Adult Asperger Assessment (AAA): a diagnostic method. *J Autism Dev Disord* 2005;35(6):807-19. Epub 2005/12/07
- 16 Lord C, Pickles A, McLennan J et al. Diagnosing autism: analyses of data from the Autism Diagnostic Interview. *J Autism Dev Disord* 1997;27(5):501-17. Epub 1997/12/24
- 17 Lord C, Risi S, Lambrecht L et al. The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *J Autism Dev Disord* 2000;30(3):205-23. Epub 2000/10/31
- 18 Gillberg C, Gillberg C, Rastam M, Wentz E. The Asperger Syndrome (and high-functioning autism) Diagnostic Interview (ASDI): a preliminary study of a new structured clinical interview. *Autism* 2001;5(1):57-66. Epub 2001/11/16

Useful information

NICE

For healthcare professionals
NICE CG142. Autism: recognition, referral, diagnosis and management of adults on the autism spectrum

For carers

NICE CG142. Diagnosing, supporting and caring for adults with autism
www.nice.org.uk

National Autistic Society
www.autism.org.uk

We welcome your feedback

If you would like to comment on this article or have a question for the authors, write to:
editor@thepractitioner.co.uk