Table 1

QOF2: Secondary prevention of coronary heart disease indicators

Indicator	Points	Payment stages
Records		
CHD 1: The practice can produce a register of patients with coronary heart disease	4	
Diagnosis and initial management		
CHD 2: The percentage of patients with newly diagnosed angina (diagnosed after 1 April 2003) who are referred for exercise testing and/or specialist assessment	7	40-90%
Ongoing management		
CHD 5: The percentage of patients with coronary heart disease whose notes have a record of blood pressure in the previous 15 months	7	40-90%
CHD 6: The percentage of patients with coronary heart disease in whom the last blood pressure reading (measured in the last 15 months) is 150/90 or less	19	40-70%
CHD 7: The percentage of patients with coronary heart disease whose notes have a record of total cholesterol in the previous 15 months	7	40-90%
CHD 8: The percentage of patients with coronary heart disease whose last measured total cholesterol (measured in the previous 15 months) is 5mmol/l or less	17	40-70%
CHD 9: The percentage of patients with coronary heart disease with a record in the previous 15 months that aspirin, an alternative antiplatelet therapy, or an anticoagulant is being taken (unless a contraindication or side-effects are recorded)	7	40-90%
CHD 10: The percentage of patients with coronary heart disease who are currently treated with a ß-blocker (unless a contraindication or side-effects are recorded)	7	40-60%
CHD 11: The percentage of patients with coronary heart disease with a history of myocardial infarction (diagnosed after 1 April 2003) who are currently treated with an ACE inhibitor or angiotensin II antagonist	7	40-80%
CHD 12: The percentage of patients with coronary heart disease who have a record of influenza immunisation in the preceding 1 September to 31 March	7	40-90%