Table 1

ILAR classification of juvenile idiopathic arthritis (JIA) categories relevant to juvenile spondyloarthropathies (JSpA)[5]

Category	Criteria
Enthesitis related arthritis (ERA)	**Diagnostic criteria** Arthritis and enthesitis **OR** Arthritis or enthesitis, with at least 2 of: ● Presence or history of sacroiliac joint tenderness and/or inflammatory lumbosacral pain ● Presence of HLA-B27 antigen ● Onset of arthritis in boys over six years of age ● Acute anterior uveitis ● History in a first-degree relative of ankylosing spondylitis, ERA, sacroiliitis with IBD, reactive arthritis, or acute anterior uveitis **Exclusion criteria** ● Psoriasis or history of psoriasis in the patient or first-degree relative ● IgM RF factor positive on two separate occasions, three months apart ● Presence of systemic JIA in the patient
Juvenile psoriatic arthritis (JPsA)	**Diagnostic criteria** Arthritis and psoriasis **OR** Arthritis with at least 2 of: ● Dactylitis ● Nail pitting or onycholysis ● History in a first-degree relative of psoriasis **Exclusion criteria** ● Arthritis in an HLA-B27 positive boy over six years old ● IgM RF factor positive on two separate occasions, three months apart ● Ankylosing spondylitis, ERA, sacroiliitis with IBD, reactive arthritis, acute, anterior uveitis or a history of one of these in a first-degree relative ● Presence of systemic JIA in the patient
Undifferentiated arthritis	Arthritis that fulfills criteria in no category, or 2 or more categories