## Table 2 Risk stratification and treatment recommendations for localised or locally

advanced prostate cancer<sup>3</sup> Cambridge Prognostic Criteria **Treatment** Group (CPG) recommendation 1 Offer active surveillance Gleason 6 (grade group 1) Consider radical and prostate specific antigen prostatectomy or radical  $(PSA) < 10 \mu g/L$ radiotherapy if active

stages T1-T2 2 Gleason score 3 + 4 = 7(grade group 2) or PSA 10-20 µg/L and stages T1-T2

and

Gleason score 3 + 4 = 7(grade group 2) and PSA 10-20 µg/L and stages T1-T2 Or Gleason 4 + 3 = 7 (grade

group 3) and stages T1-T2

One of: Gleason score 8

4

3

(grade group 4). PSA > 20 µg/L, stage T3

that the prostate cancer can be controlled in the long term Do not offer active surveillance Offer radical prostatectomy or radical

surveillance

Two or more of: Gleason score 8 (grade group 4), PSA > 20 µg/L, stage T3 Or

stage T4

radiotherapy when it is likely

prostatectomy or radical

radiotherapy when it is likely

surveillance is not suitable

Offer a choice between

active surveillance, radical

prostatectomy or radical

surveillance for people who choose not to have

prostatectomy or

Offer radical

radiotherapy

Consider active

immediate radical treatment

Offer radical

radiotherapy if radical treatment is suitable

or acceptable to the person

that the prostate cancer can

be controlled in the long term Do not offer active

5 Gleason score 9 to 10 (grade group 5) Or